

**Daniel O. Beck, MD**  
Dallas Plastic Surgery Institute  
www.beckplasticsurgery.com

---

**BRAZILIAN BUTT LIFT (GLUTEAL AUGMENTATION)  
AFTER CARE INSTRUCTIONS**

1. After surgery, it is important to have someone available to stay with you for the first 24-48 hours. You will be weak and drowsy and may require help the first few times you get out of bed.
2. While in bed lie on your stomach or side as much as possible to keep direct pressure off of the buttock region. Keep your hips flexed to 30-45 degrees with support from 2-3 pillows as necessary. Move your legs and flex/extend your ankles several times per hour. Do not cross your legs, as this increases the risk for blood clots.
3. It is very important to get out of bed (with assistance) as early and often as possible after your surgery to prevent postoperative problems. While the goal is to move about, even getting out of bed and into a chair will be beneficial. Attempt to walk 15-20 minutes 6 times per day.
4. Please take frequent deep breaths to keep your lungs clear (15-20 per hour).
5. A light diet is best after surgery. Begin by taking liquids slowly and progress to soups or Jell-O. You may start a regular diet the next day. Over the next 6 weeks, a high protein diet will benefit your recovery. Daily protein supplementation with pre-made shakes (eg. Boost, Muscle Milk) or soluble powder (whey, soy etc.) from a nutrition store is recommended.
6. Please take the pain medication as directed for the first 2-3 days after surgery. It is not uncommon for you to need it every 3-4 hours. Take it with crackers, Jell-O, etc. After this time, take the pain medication only if you have pain. If the prescription pain medication is too strong for your pain needs, over the counter ibuprofen (400-800 mg every 6-8 hours) can be substituted. In addition, you may be given an anti-anxiety/muscle relaxant medication that you may take as well to prevent aggravation of the pain cycle.
7. External sutures at the liposuction access sites and fat injection sites will dissolve and fall out on their own over the first two weeks.
8. It is normal to experience a pink fluid discharge from the liposuction access sites for several days after the procedure. This is mostly excess fluid introduced as part of the liposuction procedure. Drainage may increase as you become more mobile and stand for longer periods of time.
9. Direct pressure to the buttock must be avoided for 2 weeks after surgery to allow the transferred fat the best chance for survival. Sleeping position should be on your side or stomach. If it is absolutely necessary to sit upright, a doughnut cushion or rolled up blanket under your hips will help disperse and minimize direct pressure to the grafted site.

10. The compression garment must be worn AT ALL TIMES for 2 weeks. You may remove the garment to shower starting 72 hours after surgery. Be sure someone is with you to help, as it is not uncommon to feel dizzy or lightheaded while standing for prolonged periods of time. Sit or lie down 20 minutes prior to showering.
11. Continue to wear the compression garment at night for the 3<sup>rd</sup> and 4<sup>th</sup> week after surgery. If you wish, a full length Lycra girdle without zippers and extra support pantyhose may be purchased at a department store and substituted for the compression garment.
12. **DO NOT SMOKE** or ingest any nicotine containing products. This is very important for optimal wound healing.
13. You may gradually resume normal daily activities after 48 hours being careful to avoid any activity that causes pain or discomfort. Strenuous activities and exercises are to be avoided until 3 weeks after surgery. During this recovery period your heart rate should stay below 100 beats per minute and grand sweeping movements of the extremities avoided.
14. Prolonged soaking of the incision sites in a bathtub, hot tub, swimming pool or other body of water should be avoided for 3-4 weeks to allow proper healing to occur.
15. All incisions will be extremely sensitive to sunlight during the healing phase. Direct sun contact or tanning booths are to be avoided. Use of a clinical grade sunscreen with SPF 30 and zinc oxide (UVA and UVB protection) is recommended for at least 6 months.
16. Beginning at 3 weeks after surgery, apply a scar cream to the access incisions to maximize wound healing. Use the cream for 6 months after surgery. The addition of a hydrating cream and vitamin cream will promote early softening and maturation of these areas.

### **WHAT TO EXPECT AFTER SURGERY**

Most people experience muscle soreness as opposed to sharp pains for about 7 days after surgery. In addition, sensations like numbness, itching and burning are common during the healing process. These sensations may last several weeks and will gradually disappear.

Bruising and swelling are to be expected. Bruising can be apparent for as long as 3-4 months afterwards. The bruising will move down your body as they are absorbed.

The body retains fluid in response to surgery so do not expect to see any immediate weight loss. In addition, you may find that your clothes do not fit as easily as before for the first 3-4 weeks. Be patient. The swelling will gradually subside. In the long run, you will see more of a change in the way your clothes fit than a decrease in pounds.

Discomfort from gas and bloating can be magnified after surgery. Suggestions to minimize symptoms include limiting spicy foods, avoiding carbonated drinks, and restricting intake of sugar-free products containing Xylitol.

## **SPECIAL CONSIDERATION ON DIET & EXERCISE**

Once fat cells are collected (by liposuction) and transferred to the buttock, their survival (and ultimately your final result) is highly dependent on a favorable environment allowing nutrients and blood supply to the new fat cells. The transferred fat cells require a higher than normal level of nutrition and blood flow (your baseline metabolic rate is increased). During recovery, the failure to eat a proper diet or strenuous fat burning exercises will deplete cells of vital nutrition and decrease fat cell survival.

1. Eat a high calorie, high protein diet taking in 300-500 calories over your recommended daily intake.
2. Avoid fat burning exercises, especially those targeting the buttock, for at least 3 months following your procedure.
3. Do not attempt to lose weight for 3 months after surgery.

Once the fat cells have survived the transfer and integrated into their new environment, they will act just like the your natural buttock fat cells meaning they have the ability to grow and shrink with fluctuations in your weight.

## **MASSAGE THERAPY**

Lymphatic massage will help increase circulation and alleviate any hardness or lumpiness felt underneath the skin at the liposuction sites. Massage can begin 2 weeks after surgery. Make sure the massage therapist is well versed in lymphatic drainage techniques. 45-60 minutes of therapy is recommended 2-3 times per week for 4 weeks.

### **Please note:**

- **If you have nausea, vomiting, rash, shortness of breath, or diarrhea after taking your medications, please call our office.**
- **If you develop a fever (oral temperature greater than 100°), redness and/or increased pain at the surgical incision sites, please call Dr. Beck immediately.**

## **FOLLOW-UP INSTRUCTIONS**

It is important to be seen by Dr. Beck after your initial post-op checks. He will see you for follow-up appointments at 1 week, 3 weeks, 6 weeks, 3 months, 6 months and 1-year post-op. For your convenience you may schedule your appointments at either our Southlake or Dallas office (see below). To schedule your appointment please call us at 214-818-4751 between the hours of 8:30 am – 5:00 pm at least 1 week prior to your desired date.

## **IF YOU HAVE ANY QUESTIONS, PLEASE CALL DR. BECK**

If you need immediate care, please call Dr. Beck anytime on his cell. There is no such thing as a minor question. I would rather you call me than wonder and not do the right thing, so please call my office, or email me anytime.

Dr. Beck's cell: given at time of surgery

Dr. Beck's office: 214-818-4751

Dr. Beck's email: [drbeck@dpsi.org](mailto:drbeck@dpsi.org)

Dallas Plastic Surgery Institute- Southlake  
1601 Lancaster Drive, Suite 100  
Grapevine, TX 76051  
214-818-4751

Dallas Plastic Surgery Institute- Dallas  
9101 N. Central Expressway, Suite 600  
Dallas, TX 75231  
214-818-4751