

## COMPOUND TOPICAL ANESTHETIC INFORMED CONSENT

1. Compounded anesthetics should be used only under direct Medical Injector and Medical Aesthetician supervision in a clinical setting.
2. Avoid eye contact.
3. Avoid applying medication to large areas to prevent significant systematic adverse reactions.
4. Medical Injector and Medical Aesthetician should limit the surface area site application and use small quantities of medication to avoid potential complications.
5. The duration and frequency with which the medication is applied should be limited to avoid toxic absorption over time.
6. Clients should also inform treating Medical Injector and Medical Aesthetician of all underlying health conditions and medication history to avoid possible side effects.
7. Application site should not be covered with bandage or other tightly binding materials.
8. Pregnant or breastfeeding women should not use this medication.
9. Medication should be applied to intact skin and should not be applied to skin that is raw, blistered, burned, or has open wounds.
10. Avoid applying medication to eyes, inside nostrils, mouth, or ears or other mucus membranes.
11. Do not use medication simultaneously with other topical anesthetics.
12. Do not use medication with clients who are sensitive to para aminobenzoic acid (PABA) or are allergic to preservatives, dyes, benzocaine, lidocaine, tetracaine, dibucaine, or prilocaine.
13. Medical conditions including but not limited to blood disorders, enzyme deficiencies, liver disorders, heart disease, seizures, and allergic reactions should be evaluated by physician prior to the use of this medication.
14. Use of medication may result in the blocking of all sensation in the treated area. Therefore scratching, rubbing, and exposure to sunlight or extreme hot or cold temperatures should be avoided.
15. Use of medication may result in temporary skin discoloration or mild swelling at the application site.
16. Discontinue use of medication and seek immediate medical attention if rashes or hives appear on any part of the body, rapid and severe swelling occurs at application site or elsewhere, or if client experiences shortness of breath or difficulty breathing, drop in heart rate or blood pressure, chest pains, lightheadedness, dizziness, drowsiness or confusion, ringing or buzzing in the ears, blurred or double vision, vomiting or convulsions.
17. Do not use medication with children, the elderly, and clients with heart disorders or liver disease.
18. Do not drive or operate heavy machinery when using this medication.

**Please sign to acknowledge you read and understand the information listed above:**

Client Signature \_\_\_\_\_ Date \_\_\_\_\_