

KAROL A. GUTOWSKI, MD, FACS

AESTHETIC SURGERY

CERTIFIED BY THE AMERICAN BOARD OF PLASTIC SURGERY

MEMBER AMERICAN SOCIETY OF PLASTIC SURGEONS

AFTER CARE INSTRUCTIONS for BLEPHAROPLASTY (Eyelid Surgery)

MEDICATIONS

- Before your procedure, you will need to purchase two eye products from your pharmacy: Eye lubricant drops, and eye lubricant ointment. A good choice is **Artificial Tears Lubricant eye drops** and **Artificial Tears Lubricant eye ointment**, but other brands may be used also.
- You will be prescribed narcotic pain medication which you should take as needed and directed. While most of these medications are prescribed to be taken 1 to 2 pills every 4 to 6 hours, you may find it better to take 1 pill every 2 to 3 hours if the effects start wearing off too soon. However, do not exceed the prescribed amount per day.
- As your pain decreases, you may switch to over the counter Tylenol (acetaminophen). Keep in mind that you should not take more than 4000 mg of acetaminophen per a day. Your narcotic medication may also contain acetaminophen so you need to be aware of your total dose from all the medications you are using.
- Ibuprofen (ie-Advil, Motrin) or naproxen (Aleve) may be started the same day as your surgery and may be used together with your narcotic pain medications and with acetaminophen. However, do not use ibuprofen or naproxen if you have had problems with stomach ulcers, kidney problems or if you have been told by a doctor not to use nonsteroidal anti-inflammatory drugs (NSAIDs).
- Purchase an over the counter stool softener (Metamucil, Colace) while taking narcotic pain medication. This helps avoid constipation.

DIET

- You should resume your normal diet gradually.

ACTIVITY

- **IMPORTANT: To decrease the risk of blood clots after surgery, do not spend time in bed except for when you normally sleep. Except during your normal sleeping time, get up and walk around for at least 5 minutes every hour.**
- **SHOWER:** You may shower 1 day after surgery. DO NOT take a tub bath, swim, or whirlpool for 3 weeks or if there are any openings in the incision.
- **SLEEPING:** For the 3 days, sleep with your head & shoulders slightly elevated to help reduce swelling. Use at least 2 pillows for support.
- **Avoid bending over for the first week after surgery**
- Do **NOT** push, pull, or lift anything heavier than 20 pounds (about the weight of a 2 gallons of milk) for 1 week.
- Do **NOT** do any strenuous activity or exercise (running, weight lifting, aerobics) for 2 weeks. After you start exercising, go slowly over 1 to 2 weeks until you are comfortable with the exercises.
- Do **NOT** engage in any sexual activity for at least 2 weeks; resume when completely comfortable.
- Do **NOT** drive or handle heavy machinery while taking narcotic pain medication or muscle relaxant medications.

- Do **NOT** drive until you feel comfortable that you are no longer experiencing “blurry” vision.
- Contacts may be worn when your eyes start feeling normal and the majority of the swelling has gone away, usually about one week after surgery. If it is necessary to pull down or up on your eyelid to insert them, they should not be worn for 10 days. Glasses can be worn until then.
- Cosmetics may be worn as early as 3 days after surgery. However, eyelid or eyelash cosmetics should not be worn until 2 days after all eyelid sutures have been removed.

INCISION & SCAR MANAGEMENT

- Apply a thin layer of Vaseline ointment or antibiotic ointment to the incisions with a Q-tip 3 times a day for 3 days only. If you have tape on your incision, do **NOT** apply anything.
- Place 1 to 2 drops of the lubricating eye **drops** into each eye every 2 to 4 hours for the first 2 days or if your eyes feel dry.
- Place the lubricating eye **ointment** into each eye (pull down your lower eyelid gently and place a thin strip between the lower eyelid and your eyeball) before going to sleep for the first 2 days, or if your eyelid does not close completely, or if your eyes feel dry.
- Apply a cool pack (not ice) to the eyelids as much as possible for the first 2 days. You may purchase gel cooling pack at a pharmacy or use moist hand towels stored in a refrigerator.
- See Incision and Scar Care instructions for scar prevention. Silicone gel ointment is recommended for upper eyelid incisions after the sutures are removed. Lower eyelid incisions typically do not need scar treatment.

WHAT ELSE TO EXPECT

- You can expect some minor bleeding from the incisions and some swelling of the eyelids. If the swelling on one side is **definitely** more pronounced than on the other side, if you have a continuous bloody drip or if you are experiencing pain which is not relieved by pain medication, call Dr. Gutowski immediately.
- Tightness of the eyelid is normal after surgery. This may make it hard for you to close your eyelids completely. Your eyelids will relax with time.
- You may have some discomfort for 1 to 2 weeks, although it should gradually get better after the first 2 to 3 days.
- If there are any sutures, they will be removed in 5 to 7 days after your procedure.
- Incisions are usually thin at first, become somewhat thick and red for up to 6 months, and then begin to fade. It usually takes up to 1 year before final scar quality is established.

WHEN TO CALL DR. GUTOWSKI

- Fever over 101.5 for 2 readings taken 4 hours apart.
- Any vision changes (some blurry vision is expected for a few days).
- Marked increase in redness, swelling, or pain around incision or drains.
- Any excessive bleeding or drainage from your incisions.
- Pain is not relieved by prescription medication.
- Persistent problems with nausea or vomiting
- **CHEST PAIN or TROUBLE BREATHING: CALL 911 or go to an Emergency Room**