

**STOP ALL ASPIRIN AND BLOOD THINNERS 7-14 DAYS PRIOR TO SURGERY!!!**

\*\*\*Plavix, Coumadin, Warfarin, Aspirin, Miradon, Aggrenox, Arixtra, Persantine, Lovenox, Heparin, Ticlid, Eliquis and Xarelto all require written clearance from prescribing physician prior to stopping\*\*\*

Please do not take the following medications for 7-14 days prior to surgery:

Advil	Anaprox (Naproxen)	Azulfidine
Arthrotec (Diclofenac)	Medipren (Ibuprofen)	Naprosyn
Nuprin	Naprosyn	Motrin
Mecolmen	Ansaid (Flurbiprofen)	Aleve
Ansaid (Flurbiprofen)	Cataflam (Diclofenac Potassium)	Orudis (Ketoprofen)
Oruvail (Ketoprofen)	Celbrex	Clinoril (Sulindac)
Relafen (Nabumetone)	Day Pro (Oxaprozin)	Arava
Talacen (Pentazocine Hydrochloride)	Dolobid (Diflunisal)	Ecotrin
Tolectin (Tolmetin Sodium)	Feldene (Piroxicam)	Disalcid
Toradol (Ketorolac Tromethamine)	Indocin (Indomethacin)	Voltaren (Diclofenac Sodium)
Lodine (Etodolac)	Mobic (Meloxicam)	Methotrxat
Vicoprefen		

---

Alka-Seltzer Effervescent Tablets	Alka-Seltzer Plus Cold Medicine	Anacin Tablets & Capsules
Anodynos Tablets	Arthralgen Tablets	Arthritis Pain Tablets
Arthritis Bufferin Tablets	Arthropan Liquid	Aspirin (ASA)
Aspirin Enseals (ASA)	Ascriptin Tablets	Ascriptin A/D Tablets
Ascriptin Extra-Strength Tablets	Asperbuf Tablets	Aspergum
Bayer Aspirin (All types)	BC Tablet and Powder	Buffaprin
Bufferin (All Types)	Buffets II Tablets	Buffinol Tablets
Buf- Tablets	Cama Arthritis Pain Reliever	Cope Tablets
CP-2 Tablets	Dasin Capsules	Doan's Pills
Duoprin Capsules/Syrup	Duradyne Tablets	Ecotrim Tablets
Emagrin Tablets	Excedrin	4 Way Cold Medication
Measurin Tablets	Midol	Mobigesic Tablets
Momentum Tablets	Neocylate Tablets	Pabalate
Pepto-Bismol Tablet	Persistin Tablets	S-A-C Tablets
St. Joseph Aspirin/Cold	Saleto Tablets	Salocol
Stanback Powder	Supac	Synalgos Capsules
Trigesic	Uracel	Vanquish Caplets
Verin	Dietary/Herbal Supplements	<b>ALL VITAMINS</b>

\*\*NOTE: Following a spinal fusion **DO NOT** resume any Aspirin or NSAID products until directed by your surgeon. \*\*